

Mutations in the Fukutin-Related Protein Gene (*FKRP*) Cause a Form of Congenital Muscular Dystrophy with Secondary Laminin α 2 Deficiency and Abnormal Glycosylation of α -Dystroglycan

Martin Brockington,^{1,*} Derek J. Blake,^{3,*} Paola Prandini,¹ Susan C. Brown,¹ Silvia Torelli,^{1,4} Matthew A. Benson,³ Chris P. Ponting,² Brigitte Estournet,⁵ Norma B. Romero,⁶ Eugenio Mercuri,¹ Thomas Voit,⁷ Caroline A. Sewry,^{1,8} Pascale Guicheney,⁶ and Francesco Muntoni¹

¹The Dubowitz Neuromuscular Centre, Department of Paediatrics, Imperial College School of Medicine, Hammersmith Hospital Campus, London; ²MRC Functional Genetics Unit, ³Department of Human Anatomy and Genetics, University of Oxford, Oxford; ⁴Department of Cytomorphology, University of Cagliari, Cagliari, Italy; ⁵Hôpital Raymond Poincaré, Service de Pédiatrie, Réanimation Infantile et Rééducation Neuro-respiratoire, Garches, France; ⁶Inserm U 523, Institut De Myologie, Groupe Hospitalier Pitie-Salpêtrière, Paris; ⁷Department of Paediatrics and Paediatric Neurology, University of Essen, Essen, Germany; and ⁸Department of Histopathology, Robert Jones & Agnes Hunt Orthopaedic Hospital, Oswestry, United Kingdom

The congenital muscular dystrophies (CMD) are a heterogeneous group of autosomal recessive disorders presenting in infancy with muscle weakness, contractures, and dystrophic changes on skeletal-muscle biopsy. Structural brain defects, with or without mental retardation, are additional features of several CMD syndromes. Approximately 40% of patients with CMD have a primary deficiency (MDC1A) of the laminin α 2 chain of merosin (laminin-2) due to mutations in the *LAMA2* gene. In addition, a secondary deficiency of laminin α 2 is apparent in some CMD syndromes, including MDC1B, which is mapped to chromosome 1q42, and both muscle-eye-brain disease (MEB) and Fukuyama CMD (FCMD), two forms with severe brain involvement. The *FCMD* gene encodes a protein of unknown function, fukutin, though sequence analysis predicts it to be a phosphoryl-ligand transferase. Here we identify the gene for a new member of the fukutin protein family (fukutin related protein [*FKRP*]), mapping to human chromosome 19q13.3. We report the genomic organization of the *FKRP* gene and its pattern of tissue expression. Mutations in the *FKRP* gene have been identified in seven families with CMD characterized by disease onset in the first weeks of life and a severe phenotype with inability to walk, muscle hypertrophy, marked elevation of serum creatine kinase, and normal brain structure and function. Affected individuals had a secondary deficiency of laminin α 2 expression. In addition, they had both a marked decrease in immunostaining of muscle α -dystroglycan and a reduction in its molecular weight on western blot analysis. We suggest these abnormalities of α -dystroglycan are caused by its defective glycosylation and are integral to the pathology seen in MDC1C.

Introduction

The congenital muscular dystrophies (CMD) are a heterogeneous group of autosomal recessively inherited diseases, presenting at birth or within the first 6 mo of life with hypotonia, muscle weakness, and the variable appearance of contractures and characterized by dystrophic changes on skeletal-muscle biopsy (Dubowitz 1995). The differing degrees of motor developmental delay, physical disability, muscle pathology, and eleva-

tion of serum creatine kinase (CK), along with the variable presence of mental retardation and structural brain defects, underscores the heterogeneous nature of this condition. Recent epidemiological data suggest that the incidence of CMD is 4.65×10^{-5} and that its prevalence is 8×10^{-6} (Mostacciuolo et al. 1996), indicating that it is among the most frequent autosomal recessively inherited neuromuscular diseases.

One of the main objectives of the International Consortium on CMD has been to delineate and subdivide the various forms of CMD (Dubowitz 1994, 1995, 1997, 1999; Muntoni et al., in press). Seven genetically distinct entities have been identified so far. The most common form of CMD (~40% of cases) is due to a primary deficiency in the laminin α 2 chain of merosin (laminin-2), resulting from mutations in the *LAMA2* gene (MDC1A [MIM 156225]; Tome et al. 1994; Helbing-Leclerc et al. 1995). Other CMD disease genes identified so far are *ITGA7*, leading to integrin α 7 deficiency

Received July 20, 2001; accepted for publication September 14, 2001; electronically published October 8, 2001.

Address for correspondence and reprints: Dr. Francesco Muntoni, The Dubowitz Neuromuscular Centre, Department of Paediatrics, Imperial College School of Medicine, Hammersmith Hospital Campus, London, UK. E-mail: f.muntoni@ic.ac.uk

* These authors contributed equally to this work.

© 2001 by The American Society of Human Genetics. All rights reserved.
0002-9297/2001/6906-0005\$02.00

([MIM 600536]; Hayashi et al. 1998); *COL6A2*, coding for the $\alpha 2$ chain of collagen VI, in the Ullrich variant (UCMD [MIM 254090]; Camacho Vanegas et al. 2001); and *fukutin*, responsible for Fukuyama congenital muscular dystrophy (FCMD [MIM 253800], Fukuyama et al. 1981; Kobayashi et al. 1998). Other forms of CMD that have been mapped but for which the disease genes remain elusive are muscle-eye-brain disease (MEB [MIM 236670]), mapped to chromosome 1p32-34 (Cormand et al. 1999); a form with rigidity of the spine (RSMD1 [MIM 602771]), mapped to 1p35-36 (Moghadaszadeh et al. 1998); and a variant characterized by muscle hypertrophy (MDC1B [MIM 604801]), mapped to 1q42 (Muntoni et al. 1998; Brockington et al. 2000). The primary role of abnormalities of the extracellular matrix in these diseases is exemplified by the defects in genes for laminin $\alpha 2$, integrin $\alpha 7$, and collagen VI. In addition, basal lamina abnormalities have also been demonstrated in FCMD by means of electron microscopy (Ishii et al. 1997).

Merosin consists of three laminin chains, $\alpha 2\beta 1\gamma 1$, with $\alpha 2$ forming a link between α -dystroglycan and the basal lamina (Henry et al. 1999). α -dystroglycan is a heavily glycosylated peripheral-membrane component of the dystrophin-associated glycoprotein complex (DAPC), which, in addition to laminin $\alpha 2$, binds perlecan and agrin in the extracellular matrix, whereas β -dystroglycan, derived from the same gene (Holt et al. 2000), is a transmembrane protein that links to dystrophin intracellularly (Ibraghimov-Beskrovnaya et al. 1992). Dystroglycan therefore plays a pivotal role in linking the actin-associated cytoskeleton to components of the extracellular matrix (Ervasti et al. 1991; Ibraghimov-Beskrovnaya et al. 1992), and disruption of this axis is associated with several forms of muscular dystrophy (Campbell 1993).

FCMD is one of a number of CMD forms that have been shown to have a secondary reduction in the expression of laminin $\alpha 2$ (Hayashi et al. 1993). Recently, a profound depletion of skeletal and cardiac muscle α -dystroglycan has also been reported in FCMD (Hayashi et al. 2001). FCMD is common in Japan, because of a founder haplotype containing a 3.0-kb retrotransposal insertion into the 3' UTR of the *FCMD* gene (Kobayashi et al. 1998). Fukutin, a protein of unknown function, is predicted to be a secreted protein that has sequence similarities to a family of proteins involved in modifying cell-surface molecules such as glycoproteins and glycolipids (Aravind et al. 1999). Fukutin also contains a conserved DxD motif that is found in many glycosyltransferases (Aravind et al. 1999; Breton and Imberty 1999).

Although FCMD is rare outside Japan, other CMD syndromes with secondary laminin $\alpha 2$ deficiency with or without brain involvement are not, including MDC1B (Brockington et al. 2000), MEB (Haltia et al.

1997), and other, as-yet-unmapped variants (Talim et al. 2000; Villanova et al. 2000). We have used sequence analysis to identify fukutin homologues in the human genome as candidate genes for these variants of CMD.

In the present study, we report the identification of a gene encoding a new member of the fukutin-related-protein family (FKRP [EMBL accession number AJ314847]). This gene is mutated in a severe form of CMD, which we have called "MDC1C." Patients harboring mutations in this gene had a secondary deficiency of laminin $\alpha 2$ and a profound reduction in α -dystroglycan immunostaining. In addition, the molecular weight of α -dystroglycan was reduced in muscle. Together, these findings suggest that this protein is abnormally glycosylated in MDC1C and is central to the pathology seen in this disorder.

Families, Material, and Methods

Family 1

We recently reported the clinical features of the two children in this Scottish family and suggested that they are affected by a novel form of CMD (Mercuri et al. 2000). Both children presented soon after birth with hypotonia and feeding difficulties. They never acquired the ability to walk, because of severe weakness, which also affected their facial muscles. Weakness affected the arms more than the legs, with prominent wasting of the deltoids and pectoralis muscles, whereas both calf and quadriceps muscles were hypertrophied. Cognitive development, intelligence, and vision were normal, as was brain magnetic resonance imaging. Serum CK was markedly elevated (9,000–14,550 IU/L, normal values <190 IU/liter). The older sibling died suddenly at age 7 years, following an upper respiratory tract infection. The younger sibling is now 4 years old and recently had a gastrostomy because of feeding difficulties.

Family 2

This is a consanguineous family from Libya with one affected child. In addition, two children on the paternal side of the family had died in the first decade of life with a similar illness. She presented in the first few weeks of life with hypotonia and feeding difficulties, followed by motor delay. On examination at age 16 mo, she could not walk and was weaker in her arms than in her legs. She had calf hypertrophy and facial weakness. Her serum CK was 2,478 IU/liter and she had a myopathic EMG. Her intelligence was normal.

Family 3

The affected child in this family is a 6.5-year-old girl from southern England who is currently unable to stand or walk. She presented with hypotonia in the first few

weeks of life, followed by delayed motor milestones. She could stand with support between the ages of 4 and 6 years. Her intelligence was normal, her serum CK was markedly elevated, and she had a muscle biopsy at age 9 mo that was dystrophic. At age 6 years, she had marked hypertrophy of the leg muscles with wasting and weakness of shoulder-girdle muscles, particularly her deltoid, pectoralis, and sternocleidomastoid muscles. Serum CK was elevated (2,695 IU/liter). A brain MRI showed normal results; an echocardiogram showed mildly impaired left-ventricular function.

Family 4

The third child of this Sudanese consanguineous family is affected. A detailed clinical description of this patient and those from families 5, 6, and 7 has recently been submitted elsewhere (S. Quijano-Roy, L. Galan, A. Ferreiro, F. Cheliout-Heraut, F. Gray, M. Fardeau, A. Barois, P. Guicheney, N. B. Romero, and B. Estournet, unpublished data). He presented at birth with hypotonia, and his motor development was severely delayed; he never acquired the ability to walk. Calf and tongue hypertrophy were noticed at the ages of 4 and 10 years, respectively. At age 14 years, he went into respiratory failure and has been ventilated via a tracheostomy ever since. His cognitive development and brain MRI were normal; he developed a transient cardiac insufficiency after a respiratory infection at age 11 years, but current cardiac function is normal. Serum CK was elevated (2,277 IU/liter) at age 10 years. He underwent three muscle biopsies (at the ages of 2, 3, and 9 years) that showed a dystrophic picture with normal expression of dystrophin and sarcoglycans.

Laminin $\alpha 2$ expression was reduced with an antibody against the 80-kDa fragment and was absent with an antibody against the 300-kDa fragment. No muscle was available to study the expression of α -dystroglycan.

Family 5

The only child from this nonconsanguineous family of Caribbean origin is affected. She was symptomatic at birth with weakness and hypotonia, and her motor development was severely delayed. She could walk only with assistance between the ages of 3 and 6 years. Hypertrophy of the calves had been noticed in infancy, and macroglossia appeared around puberty. She developed progressive contractures in the elbows, knees, and fingers. Her cardiac function was normal; however, she developed restrictive respiratory insufficiency leading to nasal ventilation at age 14 years and tracheostomy at age 17 years. Spinal surgery was performed at age 16 years because of progressive scoliosis. Her serum CK was 7,760 IU/liter at age 5 years. She underwent three muscle biopsies (at the ages of 14 mo, 8 years, and 19 years) that showed a dystrophic picture with normal

expression of dystrophin and sarcoglycans. Laminin $\alpha 2$ expression was reduced with an antibody against the 80-kDa fragment and was almost absent with an antibody against the 300-kDa fragment. α -dystroglycan expression was virtually absent in her muscle. Her brain MRI and mental function were normal.

Family 6

This is a 25-year-old patient, who was the first child of a nonconsanguineous French couple. She presented at birth with hypotonia, followed by delayed motor milestones. She could stand with support between the ages of 17 mo and 7 years; hypertrophy of her calves was noticed at age 6 mo, and hypertrophy of her tongue was noticed in her second decade of life. This eventually required a partial glossectomy at age 23 years because of progressive difficulties in chewing and swallowing. Intelligence was normal. A serum CK at age 2 years was markedly elevated. A brain MRI was normal, but a mild impairment of her cardiac left-ventricular function was documented on echocardiography at age 14 years. She has been ventilated with a tracheostomy since age 13 years. A muscle biopsy showed a reduction of laminin $\alpha 2$ with an antibody that recognizes the 80-kDa fragment. No muscle was available to study the expression of α -dystroglycan.

Family 7

The patient in this French family died at age 10 years because of respiratory complications. She presented at birth with hypotonia and delayed motor milestones; she never acquired the ability to walk. Marked hypertrophy of the calves and tongue was noticed in the first years of life. Progressive feeding difficulties led to a gastrotomy at age 9 years; her intelligence and brain MRI were normal, but echocardiography showed that her left-ventricular function was at the lower limit of normal at age 9 years. She died shortly afterward because of respiratory complications. Laminin $\alpha 2$ was reduced with an antibody that recognizes the 80-kDa fragment. There was no muscle to study the expression of α -dystroglycan.

Database Screening

We performed BLAST analysis for new fukutin homologues, using the BLAST server at NCBI. The identification of transcribed sequences (ESTs) and their genomic location was also performed using BLAST analysis. Sequence alignments were obtained using the ClustalW Service at the European Bioinformatics Institute.

Northern Blot Analysis

We determined the expression of *FKRP* and an estimate of transcript size by northern blot analysis using a blot containing poly(A)⁺ RNA isolated from eight tissues (Hu-

[illegible]

b

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	52
--	---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	----

[illegible]




Figure 2 Schematic representation of the *FKRP* gene. The gene consists of 4 exons and spans no more than 12 kb. The entire coding sequence is located within exon 4. The 5' UTR is blackened, the *FKRP* open reading frame is unblackened, and the 3' UTR is hatched.

man Multiple Tissue northern blot I, Clontech). An 836-bp DNA probe containing bp 253–1088 of the *FKRP* coding sequence was amplified with primers FKRP-2F (GAGCTGGTAGACTCCTTCCT) and FKRP-4R (CCTTCTCCCATACGAAGC). The blot was hybridized with the radiolabeled (^{32}P) probe in ExpressHyb hybridization solution (Clontech), according to the manufacturer's instructions.

Radiation Hybrid Mapping

The Genebridge 4 Radiation Hybrid DNA panel (UK-MRC HGMP Resource Centre) and primers FKRP-2F and FKRP-4R, amplifying a 836-bp fragment of *FKRP*, were used to define the chromosomal localization of *FKRP*. LOD scores were calculated using the RHyME

program on the UK-MRC HGMP Resource Centre Web site.

Genotyping

Genomic DNA was extracted from whole blood by standard methods. Linkage to the *FKRP* locus was assessed by genotyping subjects for the dinucleotide repeat markers *D19S219* and *D19S606*. Genotyping was performed using an ABI 377 Automated sequencer and was analyzed using Genescan 3.1 and Genotyper 2.0 (Applied Biosystems).

Mutational Analysis

A 1.7-kb fragment containing the *FKRP* coding sequence was amplified using Advantage-GC Genomic

Figure 1 Sequence of the human *FKRP* cDNA and analysis of the primary sequence. **A**, Partial nucleotide sequence and open reading frame are shown. The 4-kb cDNA encodes a protein of 495 amino acids, with a predicted molecular weight of 54.6 kDa. **B**, Multiple sequence alignment of fukutin homologues. Missense and nonsense mutations that were found to occur within the region aligned are indicated by red asterisks (*) and slashes (/), respectively. Amino acid residues are colored according to an 80% consensus: plus signs (+) indicate positively charged residues (H, K, and R; blue), minus signs (–) indicate negatively charged residues (D and E; red), black asterisks indicate serine or threonine residues (light blue on grey background), "a" indicates aromatic residues (F, H, W, and Y; blue on yellow background), "b" indicates big residues (E, F, H, I, K, L, M, Q, R, W, and Y; blue on yellow background), "c" indicates charged residues (DEHKR; pink), "h" indicates hydrophobic residues (A, C, F, G, H, I, L, M, T, V, W, and Y; black on yellow background), "l" indicates aliphatic residues (I, L, and V; grey on yellow), "p" indicates polar residues (C, D, E, H, K, N, Q, R, S, and T; blue), and s indicates small residues (A, C, D, G, N, P, S, T, and V; green). Residues that are conserved in >80% of sequences are shown as yellow on black. Residues excised from the alignment are indicated by numbers in parentheses. Predicted secondary structures are indicated below the alignment (e/E, extended or β -strand structure; h/H, helix). Residue numbers and GeneInfo identifiers (or else cosmid names) are shown following the alignment. Ca = *Candida albicans*; Ce = *Caenorhabditis elegans*; Dm = *Drosophila melanogaster*; HELP1 = haemolysin erythrocyte lysis protein 1; Hi = *Haemophilus influenzae*; Hs = *Homo sapiens*; Ll = *Lactobacillus lactis lactis*; Pi = *Prevotella intermedia*; Rp = *Rickettsia prowazekii*; Sc = *Saccharomyces cerevisiae*; Sp = *Streptococcus pyogenes*; Tb = *Trypanosoma brucei*. Fukutin homologues partially encoded by ESTs in *T. brucei* (EST AQ655742), *T. cruzi* (AI077222), and *Leishmania major* (AQ852136), as well as a *Bacillus subtilis* sequence similar to *L. lactis* TrpF C-terminal domain (GeneInfo number 1750106; bases 12108–12419), are not shown. Two *C. elegans* sequences, marked with a dagger (†), are alternative gene predictions that differ from published versions by the prediction of additional exons.




Figure 3 Northern blot analysis demonstrating the expression of *FKRP* mRNA. A 4.0-kb transcript is seen in all tissues studied. H = heart; B = brain; Pl = placenta; Lg = lung; Lv = liver; SM = skeletal muscle; K = kidney; Pc = pancreas. It is most highly expressed in skeletal muscle, placenta, and heart and is expressed relatively weakly in the remaining tissues. An additional transcript of ~3.6 kb is visible in skeletal muscle, kidney, heart, and lung.

Polymerase Mix (Clontech) and primers FKRP-1F (AAA-GGGAATTGAGAAAGAGC) and FKRP-5 (GCTCACA-CAGAGCTTCTCC). PCR products were separated by agarose gel electrophoresis, were purified (Qiagen), and were used for direct sequencing. Sequencing reactions were performed using a ABI Prism BigDye Terminator Cycle Sequencing kit (Applied Biosystems) and primers FKRP-1R (GCAGGAAGGAGTCTACCAG), FKRP-2R (CCGAGAGGTTGAAGAGGT), FKRP-3R (CTCCTCG-TAGAGGTAGGC), FKRP-4R, and FKRP-5R. Sequencing products were separated on an ABI377 automated sequencer (Applied Biosystems) and analyzed using SeqEd (Applied Biosystems).

Immunohistochemistry

Unfixed frozen 8-mm sections were incubated with monoclonal antibodies to β -spectrin; laminins $\alpha 2$, $\beta 1$, and $\gamma 1$ (Chemicon); α - and δ -sarcoglycan (Novocastra); perlecan (Chemicon); and α -dystroglycan (V1A4-1 Upstate Biotechnology). All primary antibodies were applied for 1 h and were revealed with an appropriate biotinylated secondary antibody (Amersham 1:200) for 30 min, followed by streptavidin conjugated to Alexa (Molecular Probes) for 20 min. All dilutions and washings were made in phosphate buffered saline. Sections were mounted in aqueous mountant and were viewed with epifluorescence using a Leica Aristoplan microscope.

Immunoblotting

Muscle proteins were extracted in sample buffer consisting of 1 M Tris HCl, 1% SDS, glycerol, 2-mercaptoethanol, plus a cocktail of protease inhibitors (antipain, aprotinin, and leupeptin). Soluble protein was resolved using a NuPage Pre-cast gel (4%–12% Bis-Tris; Invitrogen) and then was transferred electrophoretically to nitrocellulose membrane (Sartorius). Nitrocellulose strips were blocked in 5% milk powder in Tris-buffered

saline buffer; were probed with antibodies to α -dystroglycan (V1A4 1:1,000), β -dystroglycan (Novocastra 1:25), and α - and γ -sarcoglycan (Novocastra 1:50, 1:100); and were washed and incubated with HRP-anti-mouse (1:50,000; Jackson). Membranes were visualized using chemiluminescence (ECL+Plus, Amersham).

Results

Database Screening

Comparison of the mouse fukutin sequence with sequence databases using TBLASTN (Altschul et al. 1997) revealed a murine expressed sequence tag (EST; AA016785) that showed significant similarity ($E = 8 \times 10^{-3}$) across the proposed fukutin active-site region. Reciprocal BLAST searches using a corrected version of this sequence as a query demonstrated its significance to fukutin homologues (Aravind and Koonin 1999), in particular *Rickettsia prowazekii* RP689 ($E = 2 \times 10^{-5}$) and putative orthologous sequences in *Homo sapiens* and *Drosophila melanogaster* (fig. 1b). The human version of this sequence was determined by a combination of EST assembly, reverse transcriptase PCR, and RACE (fig. 1a). The cDNA has a 1,488-bp open-reading frame that encodes a 495-amino-acid protein, which we have called fukutin-related protein (fig. 1a). Sequence analysis of FKRP predicts the presence of a hydrophobic transmembrane-spanning region (amino acids 4–28) followed by a “stem region” and the putative catalytic domain. A similar molecular organization is found in several Golgi-resident glycosyltransferases (Munro 1998).

Radiation-Hybrid Mapping

Radiation-hybrid mapping using the Genebridge 4 Hybridization DNA panel gave a maximum two-point LOD score of 17.25 with marker *D19S219*. The most likely order of markers is centromere-*D19S219*-*FKRP*-*D19S606*-telomere on chromosome 19q13.3. *D19S219* and *D19S606* are placed 3 cM apart on the Génethon map.

Genomic Organization and Northern Blot Analysis

The genomic organization of the *FKRP* gene was determined by comparing the cDNA sequence with a contiguous segment of chromosome 19 in contig NT-011166.2. The 12-kb gene is composed of three noncoding exons and a single large exon of 3.8-kb that contains part of the 5' untranslated region and the entire open frame and 3' untranslated region (fig. 2). Northern blot analysis demonstrated a transcript of ~4.0-kb in all tissues studied. The *FKRP* transcript is expressed predominantly in skeletal muscle, placenta, and heart and relatively weakly in the remaining tissues (fig. 3). An

additional transcript of ~3.6 kb was visible in skeletal muscle, kidney, and lung.

Mutation Analysis of the *FKRP* Gene

We initially studied the *FKRP* gene in 25 families with CMD with brain involvement. No family's results were consistent with linkage to the *FKRP* locus. In addition, direct sequencing of the *FKRP* coding region in noninformative families showed that it was normal. We concluded that *FKRP* was not a major locus for these disorders, at least in our patient population.

Next, we looked at families with CMD with a secondary deficiency in laminin $\alpha 2$ and no brain involvement. In seven families that shared a common, severe phenotype, we identified mutations in the coding region of *FKRP* that are expected to affect protein function (fig. 4 and table 1). In brief, we identified 10 different mutations—8 missense and 2 nonsense. Both affected children from consanguineous families 2 and 4 had homozygous missense changes, as did the affected child from nonconsanguineous family 5. The affected children in the remaining four families were compound heterozygotes. Families 1 and 3 had one missense and one frameshift mutation, whereas families 6 and 7 had two missense mutations (table 1). In family 1, the C1154A

change was present in the mother, but the A926G change was unexpectedly not found in the father's lymphocyte DNA, suggesting a germline mutation (fig. 4). The changes were considered to be pathogenic, because they segregated with the disease in a recessive fashion and were not present in 100 control chromosomes. A small number of silent changes were identified and are listed in table 1.

Muscle Biopsy: Immunohistochemistry and Immunoblot Analysis

The skeletal muscle biopsies showed a severe dystrophic picture in all cases (a haematoxylin- and eosin-stained section from families 1 and 2 is shown in fig. 5a–b). Antibodies to α - and β -dystroglycan in families 1 and 2 showed preserved β -dystroglycan (fig. 5e–f) but a severe depletion of α -dystroglycan on most fibers in family 1 (fig. 5c), whereas >50% of fibers showed a reduction in family 2 (fig. 5d). Only the muscle from family 1 showed reduced labeling for laminin $\alpha 2$ (fig. 5g), $\beta 1$, and $\gamma 1$, and this tended to be on the larger rather than smaller fibers. In this family, only a minority of fibers with a severe depletion of α -dystroglycan were also associated with a reduction in laminin $\alpha 2$, $\beta 1$, and $\gamma 1$ labeling. However, the muscle from family 2 did not




Figure 4 An example of mutation analysis in *FKRP*. *a*, The pedigree of family 1, showing the segregation of two heterozygous mutations, A926G and C1154A. Both affected children are compound heterozygotes, whereas the unaffected sib has inherited only one of the at-risk alleles. The father unexpectedly did not have the C1154A mutation in his lymphocyte DNA and most likely is a germline mosaic. Haplotype analysis at this and other loci confirmed paternity (data not shown). *b*, DNA sequence analysis of *FKRP*, identifying the mutations in family 1.

Table 1Mutations and Polymorphisms Found in the *FKRP* Gene

Change	Inheritance	Protein Effect
Mutations:		
Family 1 (British):		
A926G	Heterozygous	Tyr309Cys
C1154A	Heterozygous	Ser385Stop
Family 2 (Libyan):		
C1378T	Homozygous	Pro448Leu
Family 3 (British):		
G1016A	Heterozygous	Arg339His
165InsGGAG	Heterozygous	Asp60Stop
Family 4 (Sudanese):		
C649A	Homozygous	Pro217Thr
Family 5 (Caribbean):		
A1394C	Homozygous	Tyr465Ser
Family 6 (French):		
C341G	Heterozygous	Ala114Gly
G1201A	Heterozygous	Asp401Asn
Family 7 (French):		
C947G	Heterozygous	Pro316Arg
T983C	Heterozygous	Tyr328Ser
Polymorphisms:		
C-34T	Heterozygous	None
C63A	Heterozygous	None
C135T	Heterozygous	None
C249T	Heterozygous	None
G648A	Heterozygous	None
G654A	Heterozygous	None
G1341A	Heterozygous	None

show any reduction in laminin $\alpha 2$ (fig. 5h), $\beta 1$, and $\gamma 1$ labeling, suggesting that the absence of α -dystroglycan in a large proportion of fibers is not directly associated with a reduction in these laminin chains. Immunolabeling of the membrane-associated proteins β spectrin and dystrophin, along with the glycosylated proteins α - and δ -sarcoglycan, were within normal limits. Perlecan immunolabeling in families 1 and 2 was indistinguishable from controls. A marked reduction in α -dystroglycan expression was reported in two of our patients from France, the only other families from which muscle was available.

Immunoblot analysis of α -dystroglycan in families 1 and 2, adult control muscle, and Duchenne muscular dystrophy (DMD) muscle was performed. A broad band of ~156 kDa was seen in normal muscle that was reactive with antibody V1A4-1, which recognizes a glycosylated epitope (fig. 6a). Families 1 and 2 showed a narrowing of the broad reactive band with a marked shift towards a lower molecular weight. Only minor alterations in the molecular weight of α -dystroglycan were apparent in the DMD muscle, despite the presence of a similar number of immature fibers to MDC1C, as judged by the expression of neonatal myosin (not shown). Both MDC1C muscles showed the same abnormality in α -

dystroglycan on western blot. This suggests that the reduction in the mean α -dystroglycan molecular weight observed in MDC1C was not simply a reflection of an immature muscle but likely reflects differences in the pattern of glycosylation in these MDC1C patients. The reduction in β -dystroglycan in DMD was expected, given the absence of dystrophin in these patients. The molecular weight of the heavily glycosylated membrane proteins α - and γ -sarcoglycan, along with β -dystroglycan, were normal (figs. 6a–b).

Discussion

We have identified a new member of the fukutin family of proteins, FKRP, whose gene maps to chromosome 19q13.3. FKRP, like fukutin, contains sequence motifs identifying it as a putative sugar transferase (Aravind and Koonin 1999; Breton and Imberty 1999). Mutations in this gene give rise to a very distinct form of congenital muscular dystrophy, which previously has been recognized as a separate clinical entity (Mercuri et al. 2000; S. Quijano-Roy, L. Galan, A. Ferreiro, F. Cheliout-Heraut, F. Gray, M. Fardeau, A. Barois, P. Guicheney, N. B. Romero, and B. Estournet, unpublished data). The characteristic clinical features of this condition are onset in the first few weeks of life; severe weakness and wasting of the shoulder-girdle muscles; hypertrophy and weakness of the leg muscles, with inability to walk; calf and thigh hypertrophy; and severe restrictive respiratory involvement, leading to respiratory failure in the second decade of life. Several patients had signs of heart involvement, and this suggests that cardiac muscle is also affected in this disorder. Other invariable features were preserved intelligence, normal brain structure on imaging, and marked elevation of serum CK. A secondary reduction in laminin $\alpha 2$ was also observed, although this was less clear in some patients because of the severe generalized muscle damage secondary to the dystrophic muscle process. In keeping with established nomenclature, we have called this disease “MDC1C” (MDC1A represents CMD with a primary laminin $\alpha 2$ deficiency; MDC1B has a secondary laminin $\alpha 2$ deficiency and calf hypertrophy linked to 1q42). We did not observe patients with two *null* mutations in *FKRP*, a finding paralleled in FCMD, where it has been suggested that the presence of two such mutations would be embryonic lethal (Kondo-Iida et al. 1999).

Patients with MDC1C had a severe reduction in α -dystroglycan immunolabeling that was much more dramatic than the reduction in laminin $\alpha 2$, the only additional protein in which we could detect an altered pattern of expression. The muscle biopsy from family 1 showed that fibers deficient in laminin $\alpha 2$ were also deficient in α -dystroglycan immunolabeling, but the converse was not true. This suggests that whatever re-




Figure 5 Muscle sections, from families 1 (*a*) and 2 (*b*), stained with haematoxylin and eosin. Immunolabeling for α - and β -dystroglycan is shown in *c* and *e*, respectively (family 1), and in *d* and *f*, respectively (family 2). α -dystroglycan was markedly reduced on most fibers in family 1, whereas >50% of fibers showed a reduction in family 2. β -dystroglycan was normally expressed in all fibers in both families. Immunolabeling for laminin $\alpha 2$ is shown in *g* (family 1) and *h* (family 2). A reduction in laminin $\alpha 2$ expression on the larger fibers was evident only in family 1. Bar = 100 μ m.

relationship exists between the reduction in α -dystroglycan and laminin $\alpha 2$, the deficiency in α -dystroglycan occurs first. Perlecan expression was within normal limits in families 1 and 2, suggesting that other components of the basement membrane are not grossly perturbed. The normal immunolabeling of β spectrin and dystro-

phin confirmed that the alterations in α -dystroglycan were not due to generalized membrane damage, which is often a feature of dystrophic muscle. These observations, together with the reduction in the average weight of α -dystroglycan, raise the possibility that FKRP is involved in the glycosylation of α -dystroglycan.




Figure 6 a, Immunoblot analysis of muscle proteins extracted from families 1 and 2 (F1 and F2) showed a marked shift in the average molecular weight of α - but not β -dystroglycan. Control adult muscle (CTRL), together with muscle from patients with Duchenne muscular dystrophy (DMD), are included for comparison. The significant reduction in α -dystroglycan molecular weight was only observed in patients with MDC1C. b, The molecular weights of α - and γ -sarcoglycan are comparable with those of control and DMD muscle.

The normal expression of the glycosylated proteins α - and γ -sarcoglycan suggests that the pathogenesis of this disorder is closely linked to the altered expression of α -dystroglycan, rather than a more generalized effect on glycosylated membrane proteins.

Dystroglycan is widely expressed, but its glycosylation pattern varies in a tissue-specific and developmental pattern (Henry and Campbell 1999; Leschziner et al. 2000), and this may provide a possible mechanism by which defects in individual glycosyltransferases can result in a tissue-specific phenotype. Many of α -dystroglycan's binding partners, including laminin $\alpha 2$, recognize carbohydrate moieties, and their disruption would be expected to have a profound effect on its function (Durbeej et al. 1998; Montanaro et al. 1999). This is supported by the recent finding that the gene mutated in the recessive myodystrophy (*myd*) mouse encodes a glycosyltransferase, LARGE, and results in the altered glycosylation of α -dystroglycan (Grewal et al. 2001). The *myd* mouse carries an out-of-frame deletion in the *LARGE* gene that is predicted to give rise to a truncated protein lacking either of its two catalytic domains, in effect creating a functional "knockout." The *myd* phenotype is characterized by proximal muscle

weakness, reduced life span, elevated serum creatine kinase, and a skeletal-muscle pathology with foci of degeneration and regeneration. These features are shared by MDC1C; however, the *myd* mouse also shows signs of CNS involvement (sensorineural deafness) that is not apparent in MDC1C (Mathews et al. 1995). In addition, patients with MDC1C have a variable cardiac involvement not seen in the *myd* mouse. Skeletal-muscle α -dystroglycan expression in the *myd* mouse was found to be virtually absent, using the same antibody as was used in the present study. Because this antibody recognizes a glycosylated epitope, these observations suggest that an abnormality in the glycosylation of dystroglycan may underlie the skeletal-muscle pathology seen in this animal. However, no further comparisons can be made, because data were not presented on the immunohistochemical expression of either α -dystroglycan or laminin $\alpha 2$ (Grewal et al. 2001), and the depletion of α -dystroglycan on immunoblotting was more marked than that observed in MDC1C. The skeletal muscle from patients with FCMD also shows a severe immunohistochemical reduction of α -dystroglycan and almost undetectable levels of α - but not β -dystroglycan, when tested by immunoblotting. The biosynthesis of glycoproteins is a complex pathway involving the controlled activity of many different enzymes in the synthesis of a variety of carbohydrate structures. It is probable that *FKRP*, like fukutin and *LARGE*, participates somewhere along one or more of these pathways in the processing of α -dystroglycan. The more severe reduction of α -dystroglycan seen in both FCMD and the *myd* mouse, compared with that seen in MDC1C, probably represents the differing involvement of the respective glycosyltransferase in the synthesis of the glycosylated epitope recognized by the V1A4-1 antibody.

The altered processing of α -dystroglycan in MDC1C, like FCMD and the *myd* mouse, is likely to be integral to the pathology in this disease. However, although we could find no evidence of other abnormalities in the processing of other proteins, such as β -dystroglycan and α - and γ -sarcoglycan, the activities of glycosyltransferases are known to be permissive, and the existence of additional substrates and the contribution to the pathology that their altered processing would produce cannot be ignored.

The altered processing of α -dystroglycan due to its abnormal glycosylation in MDC1C, together with the recent corresponding findings in FCMD and the *myd* mouse, identifies a novel pathogenic mechanism in muscular dystrophy and opens a new avenue of research into these diseases.

Acknowledgments

We wish to thank the following organizations for financial support: the Muscular Dystrophy Campaign of Great Britain and Northern Ireland (support to F.M.), the European Community (Myo-Cluster: GENRE grant QLG1 CT 1999 00870 [to F.M.]), the Wellcome Trust (support to D.J.B.), the Association Française contre les Myopathies, and INSERM (French INSERM/AFM Research network on rare disorders) and the German Research Society (DFG) to T.V. (STR498/3-1). D.J.B. is a Wellcome Trust Senior Fellow. P.P. was supported by a University of Padua Fellowship. The expert technical assistance of Dr. Lucy Feng is also gratefully acknowledged.

Electronic-Database Information

Accession numbers and URLs for data in this article are as follows:

BLAST, <http://www.ncbi.nlm.nih.gov/BLAST/>
 ClustalW Service at the European Bioinformatics Institute, <http://www2.ebi.ac.uk/clustalw/> (for sequence alignments)
 EMBL, <http://www.ebi.ac.uk/embl/> (for human *FKRP* [accession number AJ314847])
 Online Mendelian Inheritance in Man (OMIM), <http://www.ncbi.nlm.nih.gov/Omim/> (for MDC1A [MIM 156225], MEB [MIM 236670], FCMD [MIM 253800], ITGA7 [MIM 600536], UCMD [MIM 254090], and MDC1B [MIM 604801])
 UK-MRC HGMP Resource Centre, <http://www.hgmp.mrc.ac.uk/Registered/Webapp/rhyme/index.html> (for the RHyME program)

References

- Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25:3389–3402
- Aravind L, Koonin E V (1999) The fukutin protein family—predicted enzymes modifying cell-surface molecules. *Curr Biol* 9:R836–R883
- Breton C, Imberty A (1999) Structure/function studies of glycosyltransferases. *Curr Opin Struct Biol* 9:563–571
- Brockington M, Sewry CA, Herrmann R, Naom I, Dearlove A, Rhodes M, Topaloglu H, Dubowitz V, Voit T, Muntoni F (2000) Assignment of a form of congenital muscular dystrophy with secondary merosin deficiency to chromosome 1q42. *Am J Hum Genet* 66:428–435
- Camacho Vanegas O, Bertini E, Zhang RZ, Petrini S, Minosse C, Sabatelli P, Giusti B, Chu ML, Pepe G (2001) Ullrich scleroatonic muscular dystrophy is caused by recessive mutations in collagen type VI. *Proc Natl Acad Sci USA* 98:7516–7521
- Campbell KP (1995) Three muscular dystrophies: loss of cytoskeleton-extracellular matrix linkage. *Cell* 80:675–679
- Cormand B, Avela K, Pihko H, Santavuori P, Talim B, Topaloglu H, de la Chapelle A, Lehesjoki AE (1999) Assignment of the muscle-eye-brain disease gene to 1p32-p34 by linkage analysis and homozygosity mapping. *Am J Hum Genet* 64:126–135
- Dubowitz V (1994) 22nd ENMC sponsored workshop on congenital muscular dystrophy held in Baarn, The Netherlands, 14–16 May 1993. *Neuromuscul Disord* 4:75–81
- (1995) *Muscle disorders in childhood* (2nd edition). W. B. Saunders, London
- (1997) 50th ENMC international workshop: congenital muscular dystrophy. 28 February 1997 to 2 March 1997, Naarden, The Netherlands. *Neuromuscul Disord* 7:539–547
- (1999) 68th ENMC international workshop (5th international workshop): On congenital muscular dystrophy, 9–11 April 1999, Naarden, The Netherlands. *Neuromuscul Disord* 9:446–454
- Dubowitz V, Fardeau M (1995) Proceedings of the 27th ENMC sponsored workshop on congenital muscular dystrophy. *Neuromuscul Disord* 5:253–258
- Durbecq M, Henry MD, Campbell K (1998) Dystroglycan in development and disease. *Curr Opin Cell Biol* 10:594–601
- Ervasti JM, Campbell KP (1991) Membrane organization of the dystrophin-glycoprotein complex. *Cell* 66:1121–1131
- Fukuyama Y, Osawa M, Suzuki H (1981) Congenital progressive muscular dystrophy of the Fukuyama type—clinical, genetic and pathological considerations. *Brain Dev* 3:1–29
- Grewal PK, Holzfeind PJ, Bittner RE, Hewitt JE (2001) Mutant glycosyltransferase and altered glycosylation of α -dystroglycan in the myodystrophy mouse. *Nat Genet* 28:151–154
- Haltia M, Leivo I, Somer H, Pihko H, Paetau A, Kivela T, Tarkkanen A, Tome F, Engvall E, Santavuori P (1997) Muscle-eye-brain disease: a neuropathological study. *Ann Neurol* 41:173–180
- Hayashi YK, Chou FL, Engvall E, Ogawa M, Matsuda C, Hirabayashi S, Yokochi K, Ziober BL, Kramer RH, Kaufman SJ, Ozawa E, Goto Y, Nonaka I, Tsukahara T, Wang JZ, Hoffman EP, Arahata K (1998) Mutations in the integrin $\alpha 7$ gene cause congenital myopathy. *Nat Genet* 19:94–97
- Hayashi YK, Engvall E, Arikawa-Hirasawa E, Goto K, Koga R, Nonaka I, Sugita H, Arahata K (1993) Abnormal localization of laminin subunits in muscular dystrophies. *J Neurol Sci* 119:53–64
- Hayashi YK, Ogawa M, Tagawa K, Noguchi S, Ishihara T, Nonaka I, Arahata K. (2001) Selective deficiency of α -dystroglycan in Fukuyama-type congenital muscular dystrophy. *Neurology* 57:115–121
- Helbling-Leclerc A, Zhang X, Topaloglu H, Cruaud C, Tesson F, Weissenbach J, Tome FM, Schwartz K, Fardeau M, Tryggvason K (1995) Mutations in the laminin $\alpha 2$ -chain gene (*LAMA2*) cause merosin-deficient congenital muscular dystrophy. *Nat Genet* 11:216–218
- Henry MD, Campbell KP (1999) Dystroglycan inside and out. *Curr Opin Cell Biol* 11:602–607
- Holt KH, Crosbie RH, Venzke D P, Campbell KP (2000) Biosynthesis of dystroglycan: processing of a precursor propeptide. *FEBS Lett* 468:79–83
- Kobayashi K, Nakahori Y, Miyake M, Matsumura K, Kondolida E, Nomura Y, Segawa M, Yoshioka M, Saito K, Osawa M, Hamano K, Sakakihara Y, Nonaka I, Nakagome Y, Ka-

- nazawa I, Nakamura Y, Tokunaga K, Toda T (1998) An ancient retrotransposal insertion causes Fukuyama-type congenital muscular dystrophy. *Nature* 394:388–392
- Kondo-Iida E, Kobayashi K, Watanabe M, Sasaki J, Kumagai T, Koide H, Saito K, Osawa M, Nakamura Y, Toda T (1999) Novel mutations and genotype-phenotype relationships in 107 families with Fukuyama-type congenital muscular dystrophy (FCMD). *Hum Mol Genet* 8:2303–2309
- Ibraghimov-Beskrovnya O, Ervasti JM, Leveille CJ, Slaughter CA, Sernett SW, Campbell KP (1992) Primary structure of dystrophin-associated glycoproteins linking dystrophin to the extracellular matrix. *Nature* 355:696–702
- Ishii H, Hayashi YK, Nonaka I, Arahata K (1997) Electron microscopic examination of basal lamina in Fukuyama congenital muscular dystrophy. *Neuromuscul Disord* 7:191–197
- Leschziner A, Moukhles H, Lindenbaum M, Gee SH, Butterworth J, Campbell KP, Carbonetto S (2000) Neural regulation of α -dystroglycan biosynthesis and glycosylation in skeletal muscle. *J Neurochem* 74:70–80
- Mathews KD, Rapisarda D, Bailey HL, Murray JC, Schelper RL, Smith R (1995) Phenotypic and pathologic evaluation of the *myd* mouse: a candidate model for facioscapulohumeral dystrophy. *J Neuropathol Exp Neurol* 54:601–606
- Mercuri E, Sewry CA, Brown SC, Brockington M, Jungbluth H, DeVile C, Counsell S, Manzur A, Muntoni F (2000) Congenital muscular dystrophy with secondary merosin deficiency and normal brain MRI: a novel entity? *Neuropediatrics* 31:186–189
- Moghadaszadeh B, Desguerre I, Topaloglu H, Muntoni F, Pavsek S, Sewry C, Mayer M, Fardeau M, Tome FM, Guicheney P (1998) Identification of a new locus for a peculiar form of congenital muscular dystrophy with early rigidity of the spine, on chromosome 1p35-36. *Am J Hum Genet* 62:1439–1445
- Montanaro F, Lindenbaum M, Carbonetto S (1999) α -Dystroglycan is a laminin receptor involved in extracellular matrix assembly on myotubes and muscle cell viability. *J Cell Biol* 145:1325–1340
- Mostacciuolo ML, Miorin M, Martinello F, Angelini C, Perini P, Trevisan CP (1996) Genetic epidemiology of congenital muscular dystrophy in a sample from north-east Italy. *Human Genetics* 97:277–279
- Munro S (1998) Localization of proteins to the Golgi apparatus. *Trends Cell Biol* 8:11–15
- Muntoni F, Blake D, Brockington M, Brown S, Hayashi HY, Merlini L, Topaloglu H, Sabatelli P, Voit T, Guicheney P. 6th ENMC sponsored workshop on congenital muscular dystrophy held in Nardeen, The Netherlands, 27–28 October 2000. *Neuromuscul Disord* (in press)
- Muntoni F, Taylor J, Sewry CA, Naom I, Dubowitz V (1998) An early onset muscular dystrophy with diaphragmatic involvement, early respiratory failure and secondary α 2 laminin deficiency unlinked to the *LAMA2* locus on 6q22. *Europ J Paed Neurol* 2:19–26
- Talim B, Ferreiro A, Cormand B, Vignier N, Oto A, Gogus S, Cila A, Lehesjoki AE, Pihko H, Guicheney P, Topaloglu H (2000) Merosin-deficient congenital muscular dystrophy with mental retardation and cerebellar cysts unlinked to the *LAMA2*, *FCMD* and *MEB* loci. *Neuromuscul Disord* 10:548–552
- Tome FM, Evangelista T, Leclerc A, Sunada Y, Manole E, Estournet B, Barois A, Campbell KP, Fardeau M (1994) Congenital muscular dystrophy with merosin deficiency. *C R Acad Sci III* 317:351–357
- Villanova M, Mercuri E, Bertini E, Sabatelli P, Morandi L, Mora M, Sewry C, Brockington M, Brown SC, Ferreiro A, Maraldi NM, Toda T, Guicheney P, Merlini L, Muntoni F (2000) Congenital muscular dystrophy associated with calf hypertrophy, microcephaly and severe mental retardation in three Italian families: evidence for a novel CMD syndrome. *Neuromuscul Disord* 10:541–547